



Have questions?

Need resources?

Get Help Now!

Online Help Center
<http://GetHelp.StopItNow.org>

Stop It Now![®] prevents the sexual abuse of children by mobilizing adults, families and communities to take actions that protect children before they are harmed.

Warning Signs in Children and Adolescents of Possible Child Sexual Abuse

Any one sign doesn't mean that a child was sexually abused, but the presence of several suggests that you begin asking questions and consider seeking help. Keep in mind that some of these signs can emerge at other times of stress such as:

- During a divorce
- Death of a family member or pet
- Problems at school or with friends
- Other anxiety-inducing or traumatic events

Behavior you may see in a child or adolescent

- Has nightmares or other sleep problems without an explanation
- Seems distracted or distant at odd times
- Has a sudden change in eating habits
 - Refuses to eat
 - Loses or drastically increases appetite
 - Has trouble swallowing.
- Sudden mood swings: rage, fear, insecurity or withdrawal
- Leaves "clues" that seem likely to provoke a discussion about sexual issues
- Writes, draws, plays or dreams of sexual or frightening images
- Develops new or unusual fear of certain people or places
- Refuses to talk about a secret shared with an adult or older child
- Talks about a new older friend
- Suddenly has money, toys or other gifts without reason
- Thinks of self or body as repulsive, dirty or bad
- Exhibits adult-like sexual behaviors, language and knowledge

Signs more typical of younger children

- An older child behaving like a younger child (such as bed-wetting or thumb sucking)
- Has new words for private body parts
- Resists removing clothes when appropriate times (bath, bed, toileting, diapering)
- Asks other children to behave sexually or play sexual games
- Mimics adult-like sexual behaviors with toys or stuffed animal
- Wetting and soiling accidents unrelated to toilet training

351 Pleasant Street

Suite B-319

Northampton, MA 01060

Phone: 413.587.3500

Fax: 413.587.3505

www.StopItNow.org

info@StopItNow.org

HelpLine: 1.888.PREVENT



Signs more typical in adolescents

- Self-injury (cutting, burning)
- Inadequate personal hygiene
- Drug and alcohol abuse
- Sexual promiscuity
- Running away from home
- Depression, anxiety
- Suicide attempts
- Fear of intimacy or closeness
- Compulsive eating or dieting

Physical warning signs

Physical signs of sexual abuse are rare. If you see these signs, bring your child to a doctor. Your doctor can help you understand what may be happening and test for sexually transmitted diseases.

- Pain, discoloration, bleeding or discharges in genitals, anus or mouth
- Persistent or recurring pain during urination and bowel movements
- Wetting and soiling accidents unrelated to toilet training

What You Can Do If You See Warning Signs

- Create a [Safety Plan](#). Don't wait for "proof" of child sexual abuse.
- Look for patterns of behavior that make children less safe. Keep track of behaviors that concern you. This [Sample Journal Page](#) can be a helpful tool.
- See our [Let's Talk Guidebook](#) for tips on speaking up whenever you have a concern.
- If you have questions or would like resources or guidance for responding to a specific situation, visit our [Online Help Center](#), <http://GetHelp.StopItNow.org>.

Remember, the most effective prevention takes place before there's a child victim to heal or an offender to punish.

For more information and guidance, please visit our Online Help Center, <http://GetHelp.StopItNow.org>.